

COMPLIMENTS OF YOUR IBA DEALER



For More Information Contact
Your IBA Dealer / Technician

IBA STOCK #121126

218



COMPOUND S

*New Hoof Health Products For The
Fight Against Digital Dermatitis*





IBA'S COMPOUND "S" CREAM

- IBA's Compound "S" Cream contains 40% Salicylic Acid.
- IBA's Compound "S" Cream penetrates and conditions the skin while removing the wart.
- IBA's Compound "S" Cream does not crust the skin like other gelled salicylic acid products.
- IBA's Compound "S" Cream reduces inflammation and offers almost immediate relief from the pain.
- IBA's Compound "S" Cream increases blood flow to promote healing.
- IBA's Compound "S" Cream can be applied to the wart immediately after spraying the wart with IBA's Copper Advanced and Beyond.

Indications For Use: IBA's Compound "S" Cream is indicated for use on all Bovine Heel Warts (Digital Dermatitis).

Directions For Use:

- Thoroughly clean the wart area to be treated.
- Dry the area to be treated.
- Apply enough IBA's Compound "S" Cream to the wart to thoroughly cover the wart from the base to the top and rub IBA's Compound "S" Cream into the wart area.
- If wrapping is desired, apply a wart sized dollop of IBA's Compound "S" Cream to a gauze pad and place over the wart.
- Wrap the gauze pad on the wart with IBA Flex Wrap.
- Remove the wrap after 3-5 days.
- Re-treatment may be necessary for the most advanced stubborn warts.

IBA'S COMPOUND "S" POWDER

- COMPOUND "S" Powder contains 99% SALICYLIC ACID.
- A study in The Veterinary Journal No. 198(2), published Sept 2013 and entitled: *"Efficacy Of Salicylic Acid In The Treatment Of Digital Dermatitis In Dairy Cattle"*.

Concludes as follows:

"The findings suggest Salicylic Acid should be considered as an alternative to Chlortetracycline for treatment of digital dermatitis as it appears more efficacious and would assist in reducing antibiotic use".

- As suggested in the study, 10g of Salicylic Acid powder applied topically and wrapped with a bandage is the most effective application of the powder.
- With IBA's Compound "S" Powder, we recommend 1/2 to 1 scoop (scoop included with product) of IBA's Compound "S" Powder be applied to a cotton or gauze pad and placed on the infected area and wrapped with IBA's Flex Wrap. (NOTE: 1/2 scoop equals 10g).
- Spraying the infected area with IBA's Copper Advanced & Beyond prior to applying IBA's Compound "S" Powder is highly recommended especially for advanced M3 or M4 warts.

